

<https://helda.helsinki.fi>

Convenient diastereoselective synthesis of annulated 3-substituted-(5S*,6S*,Z)-

Hassan, Alaa A.

2019-11

Hassan , A A , Mohamed , N K , Aly , A A , Tawfeek , H N , Hopf , H , Bräse , S & Nieger , M
2019 , ' Convenient diastereoselective synthesis of annulated
3-substituted-(5S*,6S*,Z)-2-(2-(2,4-dinitrophenyl)hydrazono)-5,6-diphenyl-1,3-thiazinan-4-ones
' , Molecular Diversity , vol. 23 , no. 4 , pp. 821-828 . <https://doi.org/10.1007/s11030-018-09912-5>

<http://hdl.handle.net/10138/314834>

<https://doi.org/10.1007/s11030-018-09912-5>

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.


This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Convenient Diastereoselective Synthesis of Annulated 3-substituted-(5*S**,6*S**,*Z*)-2-(2-(2,4-dinitrophenyl)hydrazono)-5,6-diphenyl-1,3-thiazinan-4-ones

Alaa A. Hassan,^{1*} Nasr K. Mohamed,¹ Ashraf A. Aly,¹ Hendawy N. Tawfeek,¹ Henning Hopf,² Stefan Bräse,³ Martin Nieger⁴



Keywords:

1,3-Thiazinan-4-ones. Substituted 2,3-diphenylacrylohydrazide. (2,4-Dinitrophenyl)-4-substituted thiosemicarbazides. 2,3-Diphenylcyclopropenone. Annulated compounds.

Racemic 2-(2,4-dinitrophenyl)hydrazono)-5,6-diphenyl-1,3-thiazinan-4-ones and (*Z*)-*N'*-(2,4-dinitrophenyl)-2,3-diphenylacrylohydrazide were formed during the diastereoselective reaction between 4-substituted 1-(2,4-dinitrophenyl)thiosemicarbazides and 2,3-diphenylcycloprop-2-enone under refluxing ethanol. The structures of the synthesized compounds were confirmed by single crystal X-ray analyses.

✉ Corresponding Authors: Alaa A. Hassan: Tel.: +0020862363011; fax: +0020862363011; e-mail: alaahassan2001@mu.edu.eg

¹ Chemistry Department, Faculty of Science, Minia University, El-Minia 61519, Egypt .

² Institut für Organische Chemie, Technische Universität Braunschweig, D-38092 Braunschweig,

³ Institute of Organic Chemistry, Karlsruhe Institute of Technology, Fritz-Haber-Weg 6, 76131 Karlsruhe, Germany.

⁴ Department of Chemistry, University of Helsinki, P.O. Box 55 (A. I. Virtasen aukio I), 00014 University of Helsinki, Finland.


1 Introduction

Thiazinanones, despite being rarely reported, are very interesting compounds due to their important role in medicinal chemistry [1-3]. Substituted thiazinanones exhibited antitumor [4], antifungal activity [5] and antimalarial activity which evaluated by Kumawat et al. [6], as well as anti-oxidant activity [7]. Reactions containing an amine, carbonyl compounds and a mercapto acid in one-pot three-component condensation or a two-step process afforded thiazinanone derivatives [5]. 3-Alkyl-2-aryl-1,3-thiazinan-4-ones containing a methylsulfonyl pharmacophore were synthesized and their cyclooxygenase-2-[COX-2] inhibitory activity have been evaluated [8]. 3-Pyridin-2-ylmethyl-1,3-thiazinan-4-ones were synthesized and their anti-oxidant activities were evaluated [7].

On the other hand, the behavior of 2,3-diphenylcyclopropanone **1** towards compounds containing C=N moieties with the formation of aza-cyclopentanones (pyrrolinones) have been reported [9-12].

Due to aza-enamine reactivity shown by the reaction of alkenylidenehydrazinecarbothioamides **2** with cyclopropanone **1** and the availability of azomethine carbon as well as sulfur atom as nucleophilic sites; 3,5-disubstituted 1,3,4-thiadiazolyl-2,3-diphenylpropanones **3** were formed [13] (Scheme 1).

The reaction of **1** with various aldehyde 4-phenylthiosemicarbazones in acetic acid provided, pyrrolo[2,1-*b*]oxadiazoles [14]. Also, thione derivatives such as 2,4-disubstituted thiosemicarbazides (**4**, R = C₆H₅; C₆H₅CH₂) reacted with **1** via a nucleophilic attack of terminal-NH₂ of **4** on the carbonyl group of **1**, afforded pyridazines **5** (Scheme 1) [15].




Scheme 1: Previously reported interaction of 2,3-diphenylcyclopropanone **1** with alkenylidene hydrazinecarbothioamides **2** and 2,4-disubstituted thiosemicarbazides **4**.

Optically active sulfur compounds play an important role in the biochemistry of many living organisms and are found in many synthetic drugs and bioactive natural products [16,17]. Recently, the [3+3] cyclization of amides with cyclopropanethiones afforded the formation of 6*H*-1,3-oxazin-6-ones and 6*H*-1,3-thiazin-6-ones [18].

Results and discussion

Herein, we report the reaction of 1-(2,4-dinitrophenyl)-4-substituted thiosemicarbazides **6a-e** with 2,3-diphenylcyclopropenone **1** in absolute ethanol under reflux, (5*S**,6*S**,*Z*)-2-(2-(2,4-dinitrophenyl)-hydrazono)-3-substituted-5,6-diphenyl-1,3-thiazinan-4-ones **7a-e** were precipitated as a major product (79 %-83%). The filtrate was subjected under chromatographic plates to give only one product namely, (*Z*)-*N'*-(2,4-dinitrophenyl)-2,3-diphenylacrylohydrazide **8** as a minor product (8%-12%) (Scheme 2).



Product	R	Yield (%)	Yield of 8 (%)
7a	PhCH ₂	81	9
7b	Ph	83	8
7c	Allyl	80	10
7d	Ethyl	79	12
7e	Cyclohexyl	80	8

Scheme 2: Preparation of 2-hydrazothiazinan-4-one derivatives **7a-e** and (*Z*)-*N'*-(2,4-dinitrophenyl)-2,3-diphenylacrylohydrazide **8**.

From the structural investigation, IR spectra of **7a-e** showed the stretching frequency range between 3265-3229 cm⁻¹ due to NH-stretching, 2935-2922 cm⁻¹ for ali-CH, 1685-1675 cm⁻¹ for C=O, 1616-1612 cm⁻¹ for C=N and 1530-1524, 1344-1335 cm⁻¹ due to nitro groups.

The ¹H NMR spectrum of **7a** (in CDCl₃) as an example showed a broad singlet at δ = 11.1 ppm due to NH-group, which was confirmed further by D₂O exchange experiment. A doublet of doublet as AX-system signals at 4.57-4.56 and 5.04-5.03 with coupling constant *J* = 4.0 Hz because of CH-6 and CH-5 of thiazinanones **7a**. The ¹H NMR spectra of **7a-e** showed the absence of any signals due to H-*N*² or H-*N*⁴ groups of **6a-e** but compound **7a** showed a doublet of doublet signals at 5.40-5.43 and 5.64-5.61 with coupling constant 15.0 Hz for diastereotopic benzyl-CH₂ group.

The ¹³C NMR spectrum of **7a** showed signals at δ = 47.15 and 47.90 ppm which were assigned to thiazinanone-CH_{6,5}. Another signals at 56.36 ppm are assigned to CH₂Ph, 168.85 ppm (C=O), 146.67 ppm

(C=N) and 144.61 ppm ((NO₂)₂-Ar-C-NH). The similarities of ¹H NMR spectra (see experimental part) reveal that the five compounds **7a-e** belong to the same gross structure type namely 3-substituted 2-(2-(2,4-dinitrophenyl)hydrazono)-5,6-diphenyl-1,3-thiazinan-4-ones. The elemental analyses and mass spectrometry of **7a-e** clearly showed that the products were formed during the addition of one molecule of **1** to one molecule of **6a-e** without any elimination.

The X-ray crystallographic structure of compound **7a** further supported its relative configurations as (rac-5*S**,6*S**,*Z*)-3-benzyl-2-(2-(2,4-dinitrophenyl)hydrazono)-5,6-diphenyl-1,3-thiazinan-4-one. The molecular of **7a** (Figure 1 and Tables 1-7, in the crystallographic data) revealed furthermore the formation of 3-benzyl-2-(2-(2,4-dinitrophenyl)hydrazono)-5,6-diphenyl-1,3-thiazinan-4-one in the *cisoid* (*Z*) structure.




Figure 1: Molecular structure of compound **7a** (displacement parameters are drawn at 50 % probability level). The crystallographic numbering does not reflect the IUPAC numbering.


The mechanism for the formation of products **7a-e** is presented in scheme 3. The sulfur atom attacks the conjugate double bond of **1** forming the intermediate **9**. Intramolecular nucleophilic attack of *N*⁴-H on C=O afforded the intermediate **11** which rearranged to give **7a-e** (Scheme 3). On the other hand, *N*⁴-H attacks the carbonyl group of **1** with the formation of **7a-e** *via* intermediate **12** (Scheme 3).

The ring opening of cyclopropanones has been reported earlier by Gomaa [19] during the reaction of *N*¹,*N*²-diarylformamidines with diphenylcyclopropanone to give 3-aryl-(*N*-4-arylformamidoyl)amino-2,3-diphenylpropionic acids.

Recently, Wu et al. reported the ring-opening acylation of cyclopropanones with organo boronic acids afforded α,β -diaryl unsaturated ketones [20].

In our study, (Z)-N'-(2,4-dinitrophenyl)-2,3-diphenylacrylohydrazide **8** was formed as a minor product (8-12%) from the reaction of **1** with **6a-e**.

The compound **8** shows IR absorption at 3320-3247 cm^{-1} due to the NH groups, strong band at 1673 cm^{-1} corresponding to carbonyl group and bands at 1528, 1344 cm^{-1} attributed to nitro groups. The ^1H NMR spectrum of **8** showed multiplet signals at 6.42 due to trisubstituted acrylohydrazide-CH, 9.43 (NH), in addition to the aromatic protons. In the ^{13}C NMR spectrum of **8**, the signal at $\delta = 165.18$ was assigned to amide-CO, 145.10 due to $(\text{NO}_2)_2\text{-Ar-C-NH}$, 133.68 and 139.75 was attributed to acrylohydrazide C2 and C3.



Scheme 3: Mechanism for the formation of 2-hydrazothiazinan-4-one derivatives **7a-e**.

The structure of (Z)-N'-(2,4-dinitrophenyl)-2,3-diphenylacrylohydrazide **8** was determined by X-ray analysis (Figure 2, Tables 8-15 in the supplementary data). The X-ray structure confirms the *trans* (*E*) geometry of the two phenyl groups with respect to the C2-C3 double bond (note that the crystallographic numbering does not correspond to the systematic IUPAC numbering rules).

The hydrazide **8** was formed *via* the nucleophilic addition of $\text{N}^2\text{-H}$ on the C=O of **1** with the formation of intermediate **13**. Elimination of RNCS from **13** afforded the formation of **8** (Scheme 4).

In order to optimize the reaction conditions, we change the solvent of the reaction to CH₃CN or CH₂Cl₂, CH₃OH, ethyl acetate and tetrahydrofuran. However, the yields of **7a-e** decreased and in some cases such as ethyl acetate and tetrahydrofuran only traces of **7a-e** were observed detectable by TLC. The excess of one of the reaction partners, namely diphenylcyclopropanone **1** or thiosemicarbazides **6a-e**, led to a significant decrease of the yields.





Figure 2: Molecular structure of **8** (minor disordered part and solvent omitted for clarity, displacement parameters are drawn at 50 % probability level).



Scheme 2: The plausible mechanism for the formation of (Z)-N'-(2,4-dinitrophenyl)-2,3-diphenylacrylohydrazide **8**.

Conclusion

Nucleophilic attack of dinitrophenyl-4-substituted thiosemicarbazides on 2,3-diphenylcyclopropenone afforded the formation of racemic 2-(2,4-dinitrophenyl)hydrazono)-5,6-diphenyl-1,3-thiazinan-4-ones as major products and (*Z*)-*N'*-(2,4-dinitrophenyl)-2,3-diphenylacrylohydrazide as minor product.

Experimental

Melting points were measured with Gallenkamp melting point apparatus. Infrared spectrum (IR) was recorded with Alpha, Bruker FT-IR instruments taken as KBr discs. ¹H NMR at 400 MHz and ¹³C NMR at 100 MHz on a Bruker AM 400 spectrometry with TMS as internal standard ($\delta = 0$), and data are reported as follows: Chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, br = broad). For ¹³C NMR, TMS ($\delta = 0$) was used as internal standard and spectra were obtained with complete proton decoupling. Mass spectra were obtained using Finnigan MAT instrument (70 eV, EI-mode). Elemental analyses for C, H, N, and S were carried out using an Elmyer 306. Preparative layer chromatography (plc) was carried out on glass plates covered with a 1.0 mm thick layer of slurry-applied silica gel (Merck Pf₂₅₄).

Starting materials

The start materials **6a-e** 2-(2,4-dinitrophenyl)-*N*-substituted hydrazinecarbothioamide were prepared from the reaction between 1-(3,5-dinitrophenyl) hydrazine and the corresponding isothiocyanates in absolute ethanol under refluxing temperature according to reported literatures [21,22]. 2,3-Diphenylcycloprop-2-enone **1** was purchased from Fluka.

General procedure:

An equimolar amounts of 2,3-diphenylcycloprop-2-enone **1** and the appropriate 1,4-disubstituted thiosemicarbazides **6a-e** were mixed in absolute ethanol and refluxed for about 4-6 h, furnished reddish orange precipitates of (5*S**,6*S**,*Z*)-2-(2-(2,4-dinitrophenyl)hydrazono)-3-substituted-5,6-diphenyl-1,3-thiazinan-4-one derivatives **7a-e** and the residue were subjected to chromatographic separation using plc and toluene/ethyl acetate (10:3) as eluent to give (*Z*)-*N'*-(2,4-dinitrophenyl)-2,3-diphenylacrylohydrazide **8** as a separated zone.

(5*S**,6*S**,*Z*)-3-benzyl-2-(2-(2,4-dinitrophenyl)hydrazono)-5,6-diphenyl-1,3-thiazinan-4-one (**7a**).

Reddish orange crystals (acetonitrile), yield 470 mg (81%), mp. 240-242 °C; IR (KBr) ν : 3265 (NH), 3072 (Ar-CH), 2928 (ali-CH), 1681 (C=O), 1612 (C=N), 1579 (Ar-C=C), 1530 and 1333 cm^{-1} (NO_2); ^1H NMR (400 MHz, CDCl_3) δ : 4.56-4.57 (dd, 1H, J = 4.0 Hz, thiazinanone-H6), 5.03-5.04 (dd, 1H, J = 4.0 Hz, thiazinanone-H5), 5.40-5.43 (d, 1H, J = 15.0 Hz, CH_2 -benzyl), 5.61-5.64 (d, 1H, J = 15.0 Hz, CH_2 -benzyl), 6.81-6.88 (m, 3H, Ar-H), 7.13-7.56 (m, 13H, Ar-H), 8.27-8.30 (m, 1H, Ar-H), 9.10 (m, 1H, Ar-H), 11.02 (br, 1H, hydrazo-NH); ^{13}C NMR (100MHz, CDCl_3) δ : 47.15 (thiazinanone-CH-6), 47.90 (thiazinanone-CH-5), 56.36 (CH_2Ph), 116.05, 123.51, 127.47, 128.01, 128.16, 128.32, 128.44, 128.67, 128.76, 129.43, 130.08, 130.25 (Ar-CH), 129.21, 132.51, 133.77, 137.05, 137.87 (Ar-C), 144.61 (Ar-C-NH), 146.67 (C=N), 168.85 (C=O); MS (m/z): 553 (M^+ , 47), 462 (13), 347 (10), 207 (51), 182 (58), 149 (56), 91 (100), 77 (15); Anal. Calcd. for $\text{C}_{29}\text{H}_{23}\text{N}_5\text{O}_5\text{S}$ (553.59): C, 62.92, H, 4.19, N, 12.65, S, 5.79. Found: C, 62.79, H, 4.06, N, 12.47, S, 5.65.

(5*S,6*S**,*Z*)-2-(2-(2,4-dinitrophenyl)hydrazono)-3,5,6-triphenyl-1,3-thiazinan-4-one (7b).**

Reddish orange crystals (acetonitrile), yield 447 mg (83%), mp. 230-232 °C; IR (KBr) ν : 3262 (NH), 3076 (Ar-CH), 2932-2922 (ali-CH), 1685 (C=O), 1615 (C=N), 1577 (Ar-C=C), 1532 and 1330 cm^{-1} (NO_2); ^1H NMR (400 MHz, CDCl_3) δ : 4.54-4.55 (d, 1H, J = 4.12 Hz, thiazinanone-H6), 5.18-5.19 (d, 1H, J = 4.12 Hz, thiazinanone-H5), 6.80-7.00 (m, 4H, Ar-H), 7.10-7.32 (m, 9H, Ar-H), 7.35-7.52 (m, 3H, Ar-H), 8.04 (m, 1H, Ar-H), 8.98 (m, 1H, Ar-H), 10.98 (br, hydrazo-NH); ^{13}C NMR (100 MHz, CDCl_3) δ : 47.60 (thiazinanone-CH5), 47.95 (thiazinanone-CH6), 115.97, 123.62, 127.71, 127.99, 128.39, 128.87, 129.35, 129.62, 130.08, 130.66, 131.03, 131.21 (Ar-CH), 129.35, 132.32, 133.77, 136.88, 137.86 (Ar-C), 144.57 (Ar-C-NH), 147.31 (C=N), 168.70 (C=O); MS (m/z): 539 (M^+ , 62), 462 (23), 357 (48), 207 (61), 135 (37), 77 (100); Anal. Calcd. for $\text{C}_{28}\text{H}_{21}\text{N}_5\text{O}_5\text{S}$ (539.56): C, 62.33; H, 3.92; N, 12.98; S, 5.94. Found: C, 62.18; H, 3.77; N, 12.87; S, 5.86.

(5*S,6*S**,*Z*)-3-allyl-2-(2-(2,4-dinitrophenyl)hydrazono)-5,6-diphenyl-1,3-thiazinan-4-one (7c).**

Reddish orange crystals (acetonitrile), yield 402 mg (80%), mp. 208-210 °C; IR (KBr) ν : 3242 (NH), 3088 (Ar-CH), 2959-2925 (ali-CH), 1682 (C=O), 1614 (C=N), 1587 (Ar-C=C), 1529 and 1342 cm^{-1} (NO_2); ^1H NMR (400 MHz, CDCl_3) δ : 4.42-4.43 (d, 1H, J = 4.08 Hz, thiazinanone-H6), 4.70-4.92 (m, 2H, allyl- CH_2N), 4.96-4.97 (d, 1H, J = 4.08 Hz, thiazinanone-H5), 5.20-5.45 (m, 2H, allyl- CH_2 =), 5.90-6.10 (m, 1H, allyl- CH =), 6.60-6.70 (m, 2H, Ar-H), 6.75-6.82 (m, 2H, Ar-H), 7.00-7.30 (m, 6H, Ar-H), 7.65 (m, 1H, Ar-H), 8.24 (m, 1H, Ar-H), 9.05 (m, 1H, Ar-H), 11.02 (br, 1H, hydrazo-NH); ^{13}C NMR (100 MHz, CDCl_3) δ :

47.05 (allyl-CH₂N), 47.15 (thiazinanone-CH₆), 47.92 (thiazinanone-CH₅), 118.87 (allyl-CH₂=), 115.99, 123.60, 128.15, 128.31, 128.41, 129.24, 130.15, 130.22, 130.68 (Ar-CH), 133.75 (allyl-CH=), 128.79, 131.89, 133.69, 137.84 (Ar-C), 144.66 (Ar-C-NH), 146.32 (C=N), 168.43 (C=O); MS (*m/z*): 503 (*M*⁺, 72), 457 (18), 404 (26), 207 (55), 182 (100), 99 (66), 77 (87); Anal. Calcd. for C₂₅H₂₁N₅O₅S (503.53): C, 59.63; H, 4.20; N, 13.91; S, 6.37. Found: C, 59.45; H, 4.07; N, 13.74; S, 6.24.

(5*S,6*S**,*Z*)-2-(2-(2,4-dinitrophenyl)hydrazono)-3-ethyl-5,6-diphenyl-1,3-thiazinan-4-one (7d).**

Reddish orange crystals (acetonitrile), yield 387 mg (79%), mp. 200-202°C; IR (KBr) *v*: 3238 (NH), 3093 (Ar-CH), 2935-2923 (ali-CH), 1678 (C=O), 1616 (C=N), 1585 (Ar-C=C), 1526 and 1330 cm⁻¹ (NO₂); ¹H NMR (400 MHz, CDCl₃) *δ*: 1.35 (t, 3H, *J* = 7.77 Hz, CH₃), 4.15-4.18 (q, 2H, *J* = 7.77 Hz, CH₂), 4.45-4.46 (d, 1H, *J* = 4.10 Hz, thiazinanone-H₆), 4.96-4.97 (d, 1H, *J* = 4.10 Hz, thiazinanone-H₅), 6.64-6.70 (m, 2H, Ar-H), 6.76-6.81 (m, 2H, Ar-H), 7.06-7.09 (m, 6H, Ar-H), 7.68 (m, 1H, Ar-H), 8.30 (m, 1H, Ar-H), 9.03 (m, 1H, Ar-H), 11.04 (br, 1H, hydrazo-NH); ¹³C NMR (100 MHz, CDCl₃) *δ*: 12.86 (CH₃), 29.06 (CH₂), 47.20, 47.70 (thiazinanone-CH_{6,5}), 116.11, 123.63, 128.00, 128.30, 128.79, 129.22, 129.78, 130.10, 130.47 (Ar-CH), 129.12, 133.43, 133.80, 134.00, 137.87 (Ar-C), 144.62 (Ar-C-NH), 146.63 (C=N), 168.60 (C=O); MS (*m/z*): 491 (*M*⁺, 48), 462 (27), 445 (35), 402 (19), 182 (100), 87 (70), 77 (91); Anal. Calcd. for C₂₄H₂₁N₅O₅S (491.52): C, 58.65; H, 4.31; N, 14.25; S, 6.52. Found: C, 58.51; H, 4.20; N, 14.07; S, 6.38.

(5*S,6*S**,*Z*)-3-cyclohexyl-2-(2-(2,4-dinitrophenyl)hydrazono)-5,6-diphenyl-1,3-thiazinan-4-one (7e).**

Reddish orange crystals (acetonitrile), yield 436 mg (80%), mp. 236-238°C; IR (KBr) *v*: 3229 (NH), 3090 (Ar-CH), 2938-2923 (ali-CH), 1675 (C=O), 1613 (C=N), 1586 (Ar-C=C), 1527 and 1331 cm⁻¹ (NO₂); ¹H NMR (400 MHz, CDCl₃) *δ*: 1.08-1.98 (m, 10H, cyclohexyl-CH₂), 2.02-2.46 (m, 1H, cyclohexyl-CH), 4.35-4.36 (d, 1H, *J* = 4.11 Hz, thiazinanone-H₆), 4.80-4.81 (d, 1H, *J* = 4.11 Hz, thiazinanone-H₅), 6.62-6.70 (m, 2H, Ar-H), 6.84-6.92 (m, 2H, Ar-H), 7.04-7.46 (m, 6H, Ar-H), 7.71 (m, 1H, Ar-H), 8.32 (m, 1H, Ar-H), 9.10 (m, 1H, Ar-H), 11.00 (br, hydrazo-NH); ¹³C NMR (100 MHz, CDCl₃) *δ*: 25.74, 26.53, 29.78 (cyclohexyl-CH₂), 47.63, 47.91 (thiazinanone-CH_{6,5}), 57.08 (cyclohexyl-CH), 115.93, 123.64, 128.04, 128.22, 128.71, 129.57, 130.25, 130.37, 130.77 (Ar-CH), 129.11, 133.38, 133.69, 137.96 (Ar-C), 144.59 (Ar-C-NH), 146.51 (C=N), 168.87 (C=O); MS (*m/z*): 545 (*M*⁺, 51), 462 (43), 499 (28), 455 (39), 182 (100), 141 (68), 77 (91); Anal. Calcd. For C₂₈H₂₇N₅O₅S (545.61): C, 61.64; H, 4.99; N, 12.84; S, 5.88. Found: C, 61.48; H, 4.83; N, 12.74; S, 5.75.

(*Z*)-*N'*-(2,4-dinitrophenyl)-2,3-diphenylacrylohydrazide (8).

Yellow crystals (acetonitrile), yield 8-12%, mp. 168-169°C; IR (KBr) ν : 3247 (NH), 3130 (Ar-H), 2930 (ali-H), 1693 (C=O), 1591 (Ar-C=C), 1542 and 1334 cm^{-1} (NO_2); ^1H NMR (400 MHz, CDCl_3) δ : 6.42 (s, 1H, acryl-CH), 7.10-7.55 (m, 10H, Ar-H), 7.94 (m, 1H, Ar-H), 8.48 (br, 1H, NH), 9.04 (m, 1H, Ar-H), 9.43 (br, 1H, amide-NH); ^{13}C NMR (100 MHz, CDCl_3) δ : 117.65, 123.18, 126.17, 127.52, 128.15, 128.26, 129.19, 129.38, 129.85 (Ar-CH), 133.68, 139.75 (C1 and C2-acrylohydrazide); 130.00, 134.34, 135.60, 137.80 (Ar-C); 145.10 (Ar-C-NH), 165.18 (C=O); MS (m/z): 404 (M^+ , 100), 356 (23), 221 (26), 205 (13), 195 (18), 181 (32), 138 (20), 77 (41); Anal. Calcd. for $\text{C}_{21}\text{H}_{16}\text{N}_4\text{O}_5$ (404.38): C, 62.37; H, 3.99; N, 13.86. Found: C, 62.25; H, 3.86; N, 13.77.

Single crystal X-ray structure determination of **7a** and **8**:

Suitable crystals were obtained by recrystallization from acetonitrile. The single-crystal X-ray diffraction study were carried out on a Bruker D8 Venture diffractometer with Photon100 detector at 123(2) K using Cu-K α radiation ($\lambda = 1.54178$ Å). Direct Methods for **7a** (SHELXS-97) [23] and dual space methods for **8** (SHELXT) [24] were used for structure solution and refinement was carried out using SHELXL-2014 (full-matrix least-squares on F^2) [25]. Hydrogen atoms were localized by difference electron density determination and refined using a riding model (H(N) free). Semi-empirical absorption corrections were applied. For **8** an extinction correction was applied. In **8** the 2,3-diphenylacrylo substituent is disordered (see cif-files for details).

Compound 7a: red crystals, $\text{C}_{29}\text{H}_{23}\text{N}_5\text{O}_5\text{S}$, $M_r = 553.58$, crystal size $0.24 \times 0.06 \times 0.02$ mm, monoclinic, space group $P2_1/c$ (No. 14), $a = 19.9611(6)$ Å, $b = 11.4148(4)$ Å, $c = 10.9667(4)$ Å, $\beta = 93.844(1)^\circ$, $V = 2493.16(15)$ Å³, $Z = 4$, $\rho = 1.475$ Mg/m³, $\mu(\text{Cu-K}\alpha) = 1.601$ mm⁻¹, $F(000) = 1152$, $2\theta_{\text{max}} = 144.2^\circ$, 21195 reflections, of which 4907 were independent ($R_{\text{int}} = 0.030$), 364 parameters, 1 restraint, $R_1 = 0.046$ (for 4478 $I > 2\sigma(I)$), $wR_2 = 0.118$ (all data), $S = 1.08$, largest diff. peak / hole = $0.723 / -0.490$ e Å⁻³.

Compound 8: yellow crystals, $\text{C}_{21}\text{H}_{16}\text{N}_4\text{O}_5 \cdot \text{C}_2\text{H}_6\text{OS}$, $M_r = 482.50$, crystal size $0.32 \times 0.16 \times 0.12$ mm, triclinic, space group $P-1$ (No. 2), $a = 9.0191(3)$ Å, $b = 11.3050(3)$ Å, $c = 11.8112(3)$ Å, $\alpha = 87.986(1)^\circ$, $\beta = 69.079(1)^\circ$, $\gamma = 79.272(1)^\circ$, $V = 1104.58(6)$ Å³, $Z = 2$, $\rho = 1.451$ Mg/m³, $\mu(\text{Cu-K}\alpha) = 1.732$ mm⁻¹, $F(000) = 504$, $2\theta_{\text{max}} = 144.0^\circ$, 15897 reflections, of which 4309 were independent ($R_{\text{int}} = 0.022$), 313 parameters, 10 restraints, $R_1 = 0.038$ (for 4234 $I > 2\sigma(I)$), $wR_2 = 0.093$ (all data), $S = 1.06$, largest diff. peak / hole = $0.584 / -0.369$ e Å⁻³.

Supporting Information

CCDC 1840977 (**7a**), and 1840978 (**8**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif

Acknowledgment

Alaa A. Hassan is indebted to the AvH-Foundation for the donation of a Shimadzu 408 IR instrument.

References

1. Rawal RK, Tripathi R, Katti SB, Pannecouque C, De Clercq E (2007) Synthesis and evaluation of 2-(2,6-dihalophenyl)-3-pyrimidinyl-1,3-thiazolidin-4-one analogues as anti-HIV-1 agents. *Bioorg. Med. Chem.* 15: 3134-3142. doi:10.1016/j.bmc.2007.02.044
2. Rawal RK, Tripathi R, Katti SB, Pannecouque C, De Clercq E (2007) Design, synthesis, and evaluation of 2-aryl-3-heteroaryl-1,3-thiazolidin-4-ones as anti-HIV agents. *Bioorg. Med. Chem.* 15: 1725-1731. doi:10.1016/j.bmc.2006.12.003
3. Verma A, Saraf SK (2008) 4-Thiazolidinone-A biologically active scaffold. *Eur. J. Med. Chem.* 43: 897-905. doi:10.1016/j.ejmech.2007.07.017
4. Kamel MM, Ali HI, Anwar MM, Mohamed NA, Soliman AM (2010) Synthesis, antitumor activity and molecular docking study of novel Sulfonamide-Schiff's bases, thiazolidinones, benzothiazinones and their C-nucleoside derivatives. *Eur. J. Med. Chem.* 45: 572-580. doi:10.1016/j.ejmech.2009.10.044
5. Verma A, Verma SS, Saraf SK (2010) A DIC mediated expeditious small library synthesis and biological activity of thiazolidin-4-one and 1,3-thiazinan-4-one derivatives. *J. Heterocycl. Chem.* 47: 1084-1089. DOI 10.1002/jhet.429
6. Kumawat MK, Singh UP, Singh B, Prakash A, Chetia D (2016) Synthesis and antimalarial activity evaluation of 3-(3-(7-chloroquinolin-4-ylamino)propyl)-1,3-thiazinan-4-one derivatives. *Arabian J. Chem.* 9: S643-S647. <http://dx.doi.org/10.1016/j.arabjc.2011.07.007>

7. Bosenbecker J, Bareño VDO, Difabio R, Vasconcellos FA, Dutra FSP, Oliveira PS, Barschak AG, Stefanello FM, Cunico WJ (2014) Synthesis and Antioxidant Activity of 3-(Pyridin-2-ylmethyl)-1,3-thiazinan(thiazolidin)-4-ones. *Biochem. Molecular Toxicology* 28: 425-432. DOI 10.1002/jbt.21581
8. Zebardast T, Zarghi A, Daraie B, Hédayati M, Dadrass OG (2009) Design and synthesis of 3-alkyl-2-aryl-1,3-thiazinan-4-one derivatives as selective cyclooxygenase (COX-2) inhibitors. *Bioorg. Med. Chem. Lett.* 19: 3162-3165. doi:10.1016/j.bmcl.2009.04.125
9. Eicher T, Franke G (1981) On the Reaction of Diphenylcyclopropanone with Guanidines. *Liebigs Ann. Chem.* 1337-1353. <https://doi.org/10.1002/jlac.198119810802>
10. Takahashi M, Funaki T, Honda H, Yokoyama Y, Takimoto H (1982) Formation of 1,2,3,5-Tetrasubstituted 2-Pyrrolin-4-ones and 1,2,3-Trisubstituted Pyrroles from Diphenylcyclopropanone and 1,4-Diazabutadienes. *Heterocycles* 19: 1921-1924. DOI: 10.3987/R-1982-10-1921
11. Gomaa MAM, Döpp D (2003) Reaction of *N*¹,*N*²-Diarylamidines with (2,3-Diphenylcyclopropan-1-ylidene)propanedinitrile: Synthesis of [2-Arylamino-4(1*H*)-pyridinylidene-[propanedinitriles. *Synthesis* 10: 1545-1548. DOI: 10.1055/s-2003-40522
12. Gomaa MAM (2002) Reaction of diimines and azines with diphenylcyclopropanone. *J. Chem. Soc.: Perkin Trans. 1*: 341-344. DOI: 10.1039/b109711n
13. Hassan AA, Abdel-Latif FF, Nour El-Din AM, Mostafa SM, Nieger M, Bräse S (2012) Synthesis of (*E*)-2,5-disubstituted 1,3,4-thiadiazolyl-2,3-diphenylpropanones from alkenylidene-hydrazinecarbothioamides. *Tetrahedron* 68: 8487-8492. <http://dx.doi.org/10.1016/j.tet.2012.07.063>
14. Aly AA, Hassan AA, Ameen MA, Brown AB (2008) Chemistry of cyclopropanones: synthesis of new pyrrolo[2,1-*b*]-1,3,4-oxadiazoles. *Tetrahedron Lett.* 49: 4060-4062. doi:10.1016/j.tetlet.2008.04.066
15. Aly AA, Hassan AA, Gomaa MAM, El-Sheref EM (2007) Unusual reactivity of thiosemicarbazides towards 2,3-diphenylcyclopropanone: synthesis of new pyridazinethiones and 1,2,4-triazolo[4,3-*b*]pyridazinethiones. *Arkivoc (xiv)*: 1-11. DOI: <http://dx.doi.org/10.3998/ark.5550190.0008.e01>
16. Clayden J, MacLellan P (2011) Asymmetric synthesis of tertiary thiols and thioethers. *Beilstein J. Org. Chem.* 7: 582-595. doi:10.3762/bjoc.7.68

- 1 17. Damani LA (1989) "Sulphur-Containing Drugs and Related Organic Compounds" Ed. Wiley:
- 2 NewYork.
- 3 18. Niu B, Jiang B, Yua LZ, Shi M (2018) Base-promoted [3 + 3] cyclization of cyclopropenones and
- 4 cyclopropenethiones with amides for the synthesis of 6H-1,3-oxazin-6-ones and 6H-1,3-thiazin-6-
- 5 ones. Org, Chem. Front. 5: 1267-1271. DOI: 10.1039/c8qo00091c
- 6 19. Gomaa, MAM (2004) Reaction of *N*¹,*N*²-Diarylamidines with 2,3-Diphenylcyclopropenone. Z.
- 7 Naturforsch, 59b: 597-600. <http://znaturforsch.com>
- 8 20. Shan L, Wu G, Liu M, Gao W, Ding J, Huang X, Wu H (2018) α,β -Diaryl unsaturated ketones via
- 9 palladiumcatalyzed ring-opening of cyclopropenones with organoboronic acids. Org, Chem. Front.
- 10 5: 1651-1654. DOI: 10.1039/c8qo00241j
- 11 21. Parashar B, Jain A, Bharadwaj S, Sharma VK (2012) Synthesis and pharmacological properties of
- 12 some novel pyrazolidine and pyrazole derivatives. Med. Chem. Res. 21:1692-1699. DOI
- 13 10.1007/s00044-011-9687-0
- 14 22. Begum S, Choudhary MI, Khan KM (2009) Synthesis, phytotoxic, cytotoxic, acetylcholinesterase
- 15 and butrylcholinesterase activities of *N,N'*-diaryl unsymmetrically substituted thioureas. Natural
- 16 Product Research 23:1719-1730. DOI: 10.1080/14786410802223778
- 17 23. Sheldrick GM (2008) A short history of SHELX. Acta Cryst. A64: 112-122.
- 18 doi:10.1107/S0108767307043930
- 19 24. Sheldrick GM (2015) SHELXT-Integrated space-group and crystal-structure determination. Acta
- 20 Cryst. A71: 3-8. doi:10.1107/S2053273314026370
- 21 25. Sheldrick GM (2015) Crystal structure refinement with SHELXL. Acta Cryst. C71: 3-8.
- 22 doi:10.1107/S2053229614024218